



Bradbury, K. J., Stuart, B., Steele, M., Yardley, L., & Little, P. (2016). Looking deeper into POWeR. *Lancet Diabetes and Endocrinology*, 4(11), 887-888. [https://doi.org/10.1016/S2213-8587\(16\)30294-7](https://doi.org/10.1016/S2213-8587(16)30294-7)

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1016/S2213-8587\(16\)30294-7](https://doi.org/10.1016/S2213-8587(16)30294-7)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Elsevier at <https://www.sciencedirect.com/science/article/abs/pii/S2213858716302947?via%3Dihub>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



The POWeR of looking into the black box


Published Online
 Junly 26, 2016
[http://dx.doi.org/10.1016/S2213-8587\(16\)30115-2](http://dx.doi.org/10.1016/S2213-8587(16)30115-2)
 See [Articles](#) page 821

Translational studies using ecologically valid components are necessary to solve real-world problems such as obesity. In *The Lancet Diabetes & Endocrinology*, Paul Little and colleagues¹ offer an example of such research in their evaluation of the Positive Online Weight Management Plus (POWeR+) intervention—a web-based behavioural weight management program. The investigators assessed the efficacy of POWeR+ with either brief face-to-face nurse support (POWeR+F) or brief remote nurse support (POWeR+R) compared with an active control strategy (evidence-based dietary advice with two brief follow-up visits). The final intervention outcomes were modest, with patients in the POWeR+F group achieving an additional weight reduction of 1.5 kg (95% CI 0.6–2.4) over 12 months, and patients in the POWeR+R group achieving an additional 1.3 kg (0.34–2.2), compared with patients in the control group. However, when considered in combination with the incremental development of the initial POWeR program,^{2,3} use of primary care facilities and behavioural theory to improve translation, and a promising economic evaluation, the intervention could prove beneficial for real-world weight management in the UK. Unfortunately, as the investigators acknowledge, the 12 month study duration does not allow for sufficient understanding of the long-term efficacy of the program. Indeed, there seemed to be early signs of weight regain based on the crude mean values presented (a 1 kg increase between months 6 and 12 in the POWeR+F group and a 1.5 kg increase in

the POWeR+R group). A similar study⁴ reported similar weight regains over a longer duration (24 months) and after higher initial losses, suggesting that there might be avenues to strengthen the POWeR+ intervention.

As have many others in the eHealth field, Little and colleagues present what is now commonly referred to as a black box—an intervention with little knowledge of its internal workings. For example, to improve its longevity, the POWeR+ intervention targets psychological factors (eg, autonomous motivation), but no evaluation of changes in these constructs is presented. Furthermore, data for receipt of the intervention offer little insight into levels of user engagement and interaction with the website—a substantial component of the intervention. The average number of interactions reported represents more than half of the intended weekly logins, and some participants failed to access core components of the website. With no description of use over time, it is impossible to ascertain whether participants accessed the site ten times over a single week, rather than receiving the intended single weekly dose of, or exposure to, content. Little and colleagues' first prerogative was to evaluate weight loss, which is a crucial, objective outcome; however, successful receipt of the intervention should also be a crucial consideration of intervention efficacy. The data presented reinforce the possible importance of this consideration. The investigators report a 2–2.5 kg difference between patients who completed more than the core website components and those who did not—a value greater than the total intervention effects. This finding adds support to previous assertions about the importance of website engagement⁵ and suggests the potential potency of website engagement for improving outcomes.

Admittedly, quantification of a person's receipt of intervention content might be difficult in real-world scenarios (eg, group education sessions). However, electronic tools and programs possess impressive ability to understand exposure, engagement, and retention, and, consequently, the intervention doses that participants receive—eg, what features were used (eg, webpages accessed), when, and for what duration? Presentation of eHealth interventions should aspire to the same rigorous standards as for other trials, and



guidelines have been developed to assist researchers working in this field.⁶ Existing analysis of usage data has led to concepts such as non-use attrition⁷ and intervention adherence⁸ that have allowed much better understanding of intervention efficacy and avenues to improve potency of web-based interventions. For example, in a large web-based trial,⁹ we witnessed weight loss results simultaneously with poor initial retention and poor uptake of specific website features. With analysis of intervention data, we were able to understand key website features that predict weight loss in order to improve future interventions. During piloting of the initial POWeR intervention, usage metrics were also analysed in great depth,¹⁰ yet no such data are reported in the present study. Not all usage data need to be explored this deeply, and simple data can still offer insight into intervention adherence, uptake, and exposure without diluting evaluation of primary outcomes or alienating specific readerships, when reported appropriately.⁴

Future studies should aspire to incorporate excellent translational strategies and rigorous research methods such as those used by Little and colleagues, but they also need to unlock the black box so interventions can rise to the challenge of long-term weight maintenance.¹¹ Some findings have suggested that technology can provide only a transient effect¹² and that face-to-face contact might be the answer to longer term maintenance of weight loss.¹³ Carefully developed, integrated interventions such as POWeR+ might be able to overcome these limitations. However, an intervention's true efficacy will only be fully understood when we can ensure that participants are receiving the intended intervention dose. This is particularly the case when

eHealth components are integral to an intervention because user engagement could be a crucial moderator of success.

Emily Brindal

CSIRO, Food and Nutrition, Adelaide, SA 5000, Australia
emily.brindal@csiro.com

I declare no competing interests

Copyright © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY license.

- 1 Little P, Stuart B, Hobbs RFD, et al. An internet-based intervention with brief nurse support to manage obesity in primary care (POWeR2): a pragmatic, parallel-group, randomised controlled trial. *Lancet Diabetes Endocrinol* 2016; published online July 26. [http://dx.doi.org/10.1016/S2213-8587\(16\)30099-7](http://dx.doi.org/10.1016/S2213-8587(16)30099-7).
- 2 Yardley L, Ware L, Smith ER, et al. Randomised controlled feasibility trial of a web-based weight management intervention with nurse support for obese patients in primary care. *Int J Behav Nutr Phys Act* 2014; **11**: 67.
- 3 Yardley L, Morrison L, Bradbury K, Muller I. The person-based approach to intervention development: application to digital health-related behavior change interventions. *J Med Internet Res* 2015; **17**: e30.
- 4 Appel LJ, Clark JM, Yeh HC, et al. Comparative effectiveness of weight-loss interventions in clinical practice. *N Engl J Med* 2011; **365**: 1959–68.
- 5 Neve M, Morgan PJ, Jones PR, Collins CE. Effectiveness of web-based interventions in achieving weight loss and weight loss maintenance in overweight and obese adults: a systematic review with meta-analysis. *Obesity Rev* 2010; **11**: 306–21.
- 6 Eysenbach G, CONSORT-EHEALTH Group. CONSORT-EHEALTH: improving and standardizing evaluation reports of web-based and mobile health interventions. *J Med Internet Res* 2011; **13**: e125.
- 7 Eysenbach G. The law of attrition. *J Med Internet Res* 2005; **7**: e11.
- 8 Kelders SM, Kok RN, Ossebaard HC, Van Gemert-Pijnen JEW. Persuasive system design does matter: a systematic review of adherence to web-based interventions. *J Med Internet Res* 2012; **14**: 17–40.
- 9 Brindal E, Freyne J, Saunders I, Berkovsky S, Smith G, Noakes M. Features predicting weight loss in overweight or obese participants in a web-based intervention: randomized trial. *J Med Internet Res* 2012; **14**: e173.
- 10 Arden-Close EJ, Smith E, Bradbury K, et al. A visualization tool to analyse usage of web-based interventions: the example of Positive Online Weight Reduction (POWeR). *JMIR Hum Factors* 2015; **2**: e8.
- 11 Wing RR, Phelan S. Long-term weight loss maintenance. *Am J Clin Nutr* 2005; **82** (1 suppl): 222S–5S.
- 12 Svetkey LP, Stevens VJ, Brantley PJ, et al. Comparison of strategies for sustaining weight loss: the weight loss maintenance randomized controlled trial. *JAMA* 2008; **299**: 1139–48.
- 13 Wing RR, Tate DF, Gorin AA, Raynor HA, Fava JL. A self-regulation program for maintenance of weight loss. *N Engl J Med* 2006; **355**: 1563–71.

Knowing what we do not know: statin therapy in advanced chronic kidney disease



Lowering LDL cholesterol with statins clearly reduces the risk of cardiovascular events among individuals at increased cardiovascular risk, and is a mainstay of contemporary approaches to cardiovascular disease prevention. Chronic kidney disease is a well established independent risk factor for cardiovascular events, but guidelines vary in their approach to lipid-lowering in

this population. The European Society of Cardiology/European Atherosclerosis Society (ESC/EAS) guidelines suggest that chronic kidney disease is an indication for statin therapy, whereas the American College of Cardiology (ACC)/American Heart Association (AHA) guidelines do not.¹ The KDIGO (Kidney Disease: Improving Global Outcomes) guidelines recommend

Published Online
July 28, 2016
[http://dx.doi.org/10.1016/S2213-8587\(16\)30192-9](http://dx.doi.org/10.1016/S2213-8587(16)30192-9)
See [Articles](#) page 829